

Quasi- Secondary Submodules

A. J. Taherizadeh: Kharazmi University

Abstract

Let R be a commutative ring with non-zero identity and M be a unitary R -module. Then the concept of quasi-secondary submodules of M is introduced and some results concerning this class of submodules is obtained.

1. Introduction

Throughout this paper all rings are commutative with non-zero identity and all modules are unitary. In [4] L.Fuchs introduced and studied the concept of quasi-primary ideals (see also [5]). An ideal I of a ring R is called a *quasi-primary* ideal of R if the radical of I is a prime ideal of R . This concept then generalized to modules, i.e., the concept of quasi-primary submodules of a module introduced and developed in [3]. Here, we introduce the dual notation, that is, the quasi-secondary submodules of a module and obtain some results concerning this class of submodules. In section 2, we obtain some preliminary properties of quasi-secondary submodules. Section 3 is devoted to the quasi-secondary submodules of a multiplication module. Now we define some concepts which will be needed in sequel.

Let M be an R -module and N a submodule of it. The ideal $\{r \in R \mid rM \subseteq N\}$ will be denoted by $(N \dot{\vdash}_R M)$; in particular $(0 \dot{\vdash}_R M)$ is called the annihilator of M . A non-zero submodule N of M is called a *secondary* (resp. *second*) submodule of M if for each $r \in R$ the homothety $N \xrightarrow{r} N$ is surjective or nilpotent (resp. surjective or zero). In this case $\sqrt{(0 \dot{\vdash}_R N)}$ is a prime ideal, say p , and we call N a *p-secondary* (resp. a *p-second*) submodule of M . We refer readers for more details concerning secondary (resp. second) submodule to [9] (resp. [12]).

KeyWords: quasi – secondary submodules, secondary submodules, multiplication modules

2010 Mathematics Subject Classification: 13C05, 13C13

Received: 26 Nov. 2011

Revised 18 July 2012

* Correspondence Author Taheri@tmu.ac.ir

An R -module M is said to be a *multiplication* module if for each submodule N of M there exists an ideal I of R such that $N = IM$. It is easy to see that in this case $N = (N_R M)M$. Also the ideal $\theta(M)$ is defined as $\theta(M) := \sum_{m \in M} (Rm_R M)$. If M is a multiplication module and N is a submodule of it, then $M = \theta(M)M$ and $N = \theta(M)N$. (see [1]). An R -module M is *sum-irreducible* if $M \neq 0$ and the sum of any two proper submodules of M is always a proper submodule. Finally a proper submodule N of an R -module M is called a *prime submodule* if for each $r \in R$ the homomethy $M/N \xrightarrow{r} M/N$ is either injective or zero. This implies that $\text{Ann}(M/N) = p$ is a prime ideal of R , and N is said to be a *p-prime submodule* (c.f. [7], [8], [10] and [11]).

2. Quasi-Secondary Submodules

The starting point of this section is the definition of quasi-secondary submodules of a module.

Definition 2.1. Let M be a non-zero R -module. Then the non-zero submodule N of M is said to be *quasi-secondary* if $\sqrt{(0_R N)} = p$ where p is a prime ideal of R . It is obvious that every secondary (or second) submodule of a module is a quasi-secondary submodule, but the converse is not true in general. For example, $2Z$ is a 0-quasi-secondary submodule of the Z -module Z but it is not 0-secondary (or 0-second) submodule. (Here Z denotes the set of all integers.)

Remark 2.2.

- (i) Let M be a non-zero R -module and N a submodule of it such that $\sqrt{(0_R N)} = m(m \in \text{Max}(R))$. Then N is *m-secondary* (*m-second*).
- (ii) Every quasi-secondary submodule of a module over a zero-dimensional ring (i.e., a ring in which every prime ideal is a maximal ideal) is secondary.
- (iii) Every quasi-secondary submodule of a module over a D.V.R is secondary.

Definition 2.3. Let M be an R -module and N a submodule of M . An element r of R is called *co-primal* to N if $rN = N$. Denote by $W(N)$ the set of all elements of R that are not co-primal to N . The submodule N is said to be a *co-primal submodule* of M if $W(N)$ is an ideal of R . This ideal is always a prime ideal. In this case we say that N is a *p-co-primal* submodule of M . The class of co-primal submodules of a module is a

fairly large class. For example , all secondary (second) submodules are co-primal. Also it is easy to see that a sum-irreducible submodule of a module is co-primal. But, in general, a quasi-secondary submodule of a module may not be a co-primal submodule. (consider the Z -module Z .) It is worth to mention that in [2] the term secondal is used for co-primal submodules. The next proposition characterizes those p -quasi- secondary submodules which are p -co-primal.

Proposition 2.4. Let N be a p -quasi-secondary submodule of an R -module M . Then N is a p -co-primal submodule of M if and only if it is a p -secondary submodule of M .

Proof \Rightarrow) Let $N \xrightarrow{r} N$ be the R -endomorphism of N given by multiplication by r of R and $rN \neq N$. Then by our assumption $r \in p = \{s \in R \mid sN \neq N\}$. On the other hand, $p = \sqrt{0_R}N$ and so there exists a positive integer t such that $r^t N = 0$. The result follows. \Leftarrow) Is obvious.

The proof of two next propositions is easy and so we state them without proof.

Proposition 2.5. Let M be a module over an integral domain and N be a 0 - co-primal submodule of M . Then N is 0 -secondary.

Proposition 2.6. Let M be an R -module and N_1, N_2, \dots, N_t be submodules of M . Then

- (i) Suppose that for $i = 1, 2, N_i$ is p_i -quasi-secondary. Then $N_1 + N_2$ is quasi-secondary if and only if $p_1 \subseteq p_2$ or $p_2 \subseteq p_1$
- (ii) If N_1, \dots, N_t are p -quasi-secondary, then $N_1 + \dots + N_t$ is a p -quasi-secondary submodule of M .
- (iii) If $N_1 + \dots + N_2$ is a p -quasi-secondary submodule of M . Then N_j is p -quasi-secondary for some $j, 1 \leq j \leq t$.

3. Multiplication Modules

In this short section we give a property of quasi-secondary submodules of a multiplication module .

Lemma 3.1. let M be a multiplication module and N be a p -quasi-secondary submodule of M . Then $\theta(M) \not\subseteq p$.

Proof. Suppose that $\theta(M) \subseteq p$ and $0 \neq n \in N$. Then $Rn = \theta(M)Rn \subseteq pn$. Hence $n = p_0n$ for some $p_0 \in p$. By our assumption there exists a positive integer t such that $p_0^t N = 0$. Therefore $n = p_0^t n = 0$, a contradiction.

Theorem 3.2. Suppose that M is a faithful multiplication module and N a p -quasi-secondary submodule of M . Then pM is a prime submodule of M . In particular, if $p \in \max(R)$, then pM is a maximal submodule of M .

Proof. By Lemma 3.1, $\theta(M) \not\subseteq p$. Now suppose that $pM = M = RM$. Then by [1, Theorem 1.5] $R \cap \theta(M) = \theta(M) = p \cap \theta(M)$ and hence $\theta(M) \subseteq p$ which is a contradiction. Thus $pM \neq M$ and the result of the first part follows from [6, Lemma 2.4(2)]. The last part can be deduced from the first part and [6, Corollary 2.7]

Acknowledgement

The author would like to thank Kharazmi University for the financial support.

References

1. Y. Al-shaniafi and S. Singh, "A companion ideal of a multiplication module", *Periodica Mathematica Hungarica.*, 46 (1) (2003) 1-8.
2. H. Ansari-Toroghy and F. Farshadifar, "The dual notions of some generalizations of prime submodules", *J. Algebra*, 39 (2011) 2396-2416.
3. S. Ebrahimi Atani and A.Y. Darani, "On quasi-primary submodules", *Chiang May J.Sci.*, 33(3) (2006) 249-254.
4. L. Fuchs, "On quasi-primary ideals", *Acta.Sci.Math. (Szeged)*, 11 (1947) 174-183.
5. L. Fuchs and E. Mosteig, "Ideals theory in prufer domains", *J. Algebra*, 252 (2002) 411-430.
6. S. C. Lee, S. Kim, S. C. Chung, "Ideals and submodules of multiplication modules", *J. Korean Math. Soc.* 42 (5) (2005) 933-948.
7. C. P. Lu, "Prime submodules of modules", *Comm. Math. Univ. Sancti. Pauli*, 33 (1984) 61-69.
8. C. P. Lu, "Spectra of modules", *Comm.in Algebra*, 23 (1995) 3741-3752.
9. I. G. Macdonald, "Secondary representation of modules over commutative rings", *Symp. Math.* XI (1973) 23-43.
10. A. Marcelo, J. Masque, "Prime submodules, the descent invariant and modules of finite length", *J.of Algebra*, 189(1997) 273-293.
11. R. L. McCasland, P.F. Smith, "Prime submodules of Noetherian modules", *Rocky Mtn. J.* 23 (1993) 1041-1062.
12. S. Yassemi, "The dual notion of prime submodules", *Arch. Math. (Brno)* 37 (2001) 273-278.