مقاله مرورى Review Article

Nova Biologica Reperta 9(2): 95-114 (2022)

Print ISSN: 2423-6330/Online ISSN: 2476-7115

https://nbr.khu.ac.ir; Kharazmi University Press; DOI: 10.29252/nbr.9.2.95

یافتههای نوین در علوم زیستی جلد ۹، شماره ۲، صفحات ۹۵ الی ۱۱۴ (۱۴۰۱) انتشارات دانشگاه خوارزمی

اثرات تغییر اقلیم و عوامل هواشناختی بر شاخصهای فصل گردهای آرایههای گیاهی آلرژیزا

فاطمه موسوی 1 ، یوسف شاه علی 7 ، خوزه اوتروس 6,7 و کارل کریستین برگمان 6

امرکز زیست فضا و محیط زیست، پژوهشگاه هوافضا، وزارت علوم، تحقیقات و فناوری، تهران، ایران، ایران، امرکز تحقیقات واکسن و سرم سازی رازی، سازمان تحقیقات، آموزش و ترویج کشاورزی، کرج، ایران، ایران، اگروه گیاه شناسی، اکولوژی و فیزیولوژی گیاهی، دانشگاه کوردوبا، اسپانیا؛ امرکز آلرژی و محیط زیست (ZAUM)، عضو مرکز تحقیقات ریه آلمان (DZL)، دانشگاه فنی مونیخ / مرکز هلمهولتز، مونیخ، آلمان ^۵ مرکز آلرژی-چاریته، دانشگاه پزشکی چاریته، برلین، آلمان به moosavi @ari.ac.ir و پوسف شاه علی، y.shahali @rvsri.ac.ir

چکیده. آلرژی های تنفسی در دو دهه گذشته تا حدی در نتیجه تغییر اقلیم در حال افزایش هستند. برای بسیاری از گونه های درختی، گراس و علف هرز افزایش تولید گرده و طول مدت گرده افشانی، منجر به افزایش طولانی مدت آلرژن های گرده ای در جو؛ جابجایی های زودهنگام دانه های گرده هوابرد و تماس طولانی مدت با آلرژن های تنفسی با اثرات مهم سلامتی بر روی افراد آلرژیک می گردد. هدف از این مقاله بررسی اثر تغییر اقلیم و عوامل هواشناختی بر روی شاخص های فصل گرده ای با یک تمرکز ویژه بر روی تاکسون های اصلی آلرژی زا در سرتاسر جهان است. متغیرهای اصلی تاثیرگذار بر فنولوژی گلدهی مانند موقعیت مکانی، عوامل اقلیمی و هواشناختی، شناسایی، مورد بحث و اثبات قرار گرفتند. دما، تشعشع خورشیدی، رطوبت، سرعت و جهت باد، از جمله مهمترین عوامل هواشناسی اثرگذار بر نوسانات غلظت سالانه گرده های هوابرد آلرژی زا هستند. اگرچه تغییرات قابل توجهی مطابق با گونه های آلرژی زا و ناحیه چغرافیایی مورد مطالعه مشاهده شد، اما به نظر می رسد دما مهمترین عامل اقلیمی اثرگذار بر فنولوژی گلدهی و تولید گرده در چندین گونه آلرژی زای های درختی باشد. افزایش سطوح دی اکسیدکربن همچنین منجر به افزایش زیست توده گیاهی، افزایش شدت گلدهی و تولید گرده در چندین گونه آلرژی زاید درختی ، گراس و علف هرز می گردد. در پرتو این بررسی شواهد فزاینده ای وجود دارد که از تاثیر تغییر اقلیم بر فنولوژی گلدهی و شاخص های فصل گرده ای تعداد قابل توجهی از گونه های گیاهی مهاجم و زینتی آلرژی زا حمایت می کند.

واژههای کلیدی. آئروبیولوژی، آلرژی، هوابرد، تغییر اقلیم، عوامل هواشناختی، فنولوژی گلدهی

The impacts of climate change and meteorological factors on pollen season indicators of allergenic plant taxa

Fateme Mousavi¹, Youcef Shahali², José Oteros^{3, 4} & Karl-Christian Bergmann⁵

Space Biology and Environment Center, Aerospace Research Institute (ARI), Ministry of Science Research and Technology (MSRT), Tehran, Iran; ² Razi Vaccine and Serum Research Institute (RVSRI); Agricultural Research, Education and Extension Organization (AREEO); Karaj, Iran; ³ Departments of Botany, Ecology and Plant Physiology, University of Córdoba (UCO), Córdoba, Spain; ⁴ Center of Allergy & Environment (ZAUM), Member of the German Center for Lung Research (DZL), TechnischeUniversität München/Helmholtz Center, Munich, Germany; ⁵ Allergy-Centre-Charité, Charité–Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universitätzu Berlin, and Berlin Institute of Health

Corresponding authors: Fateme Mousavi, moosavi@ari.ac.ir and Youcef Shahali, y.shahali@rvsri.ac.ir

Abstract. Pollen respiratory allergies have been increasing in prevalence over the last two decades, partly as the result of the impact of climate change. For many allergenic trees, grass and weed species, increased pollen production and prolonged pollination period result in long-term increased abundance of pollen allergens in the atmosphere; earlier shifts of airborne pollen grains and prolonged exposure to respiratory allergens with important health effects on allergic individuals. The aim of this review paper was to investigate the impact of climate change and meteorological factors on pollen season indicators with a special focus on the main allergenic taxa worldwide. Main variables influencing flowering

phenology such as location, climatic and meteorological parameters were identified, discussed and substantiated by published literature. Temperature, solar radiation, humidity, rainfall, wind speed and direction were identified among the most important meteorological parameters affecting the fluctuations of annual concentrations of allergenic airborne pollen grains. Although notable variations were observed according to allergenic species and studied geographical areas, temperature appeared to be the most important climatic parameter affecting flowering phenology and pollen season indicators, especially in tree species. Rising carbon dioxide levels also result in increased plant biomass, increased flowering intensity and pollen production in several tree, grass and weed allergenic species. In the light of this review, there is a growing body of evidence supporting the effect of climate change on the flowering phenology and pollen season indicators of a substantial number of allergenic ornamental and invasive plant species.

Key words. aerobiology, allergy, airborne, climate change, meteorological factors, flowering phenology

Introduction

The issue of climate change, and particularly global warming resulting from anthropogenic activities, is one of the major environmental concerns that have attracted the attention of scientists and policy makers in the last two decades. This phenomenon is considered as one of the ten main factors that can endanger the life of living organisms (Doran & Zimmerman, 2009). Climate change means stable and long-term change in the Earth's climate patterns (caused by changes in climate components and the relationships between them) which is triggered by natural events and human activities (Ghahremaninejad et al., 2021; Solomon et al., 2007).

Airborne particles are all particles that are passively transferred by air. These particles may originate from biological source (bioaerosol) or nonbiological (aerosol). Bioaerosols represent particles of biological origins with variable sizes ranging from 0.001 to 100 µm (Georgakopoulos et al., 2008). They include bacteria, viruses, pollen, fungi spores, bacterial endotoxins, antigens, allergens, toxins, mycotoxins, glucans, and plant fibers (Rogoff, 2013). The prevalence of bioaerosols in the environment is associated with many risks to human health such as pneumonia, influenza, allergies, SARS-CoV-2, etc. (Srikanth et al., 2008) Pollen grains are large biological aerosol particles with a diameter of 10-100 µm constituting the major source of respiratory allergy worldwide (Cariñanos & Casares-Porcel, 2011; Charpin et al., 2019; Sénéchal et al., 2015). Anemophilous plant species release huge quantities of pollen grains into the atmosphere that are able to interact with other atmospheric components such as air pollutants, impacting their allergenic properties (Chassard et al., 2015; Duque et al., 2013; Frank & Ernst, 2016; Okuyama et al., 2007; Sénéchal et al., 2015). Aerobiology is the science studying airborne biological particles including pollen; it also examines the impact of climate modifications on bioaerosols. Aeropalynology specifically examines the behavior, interactions and biology of pollen

grains released in the atmosphere (Cecchi et al., 2010; Corden & Millington, 2001).

Many airborne pollen grains released by tree, grass and weed species can cause pollinosis in susceptible individuals with symptoms such as watery eyes, eye irritation, runny nose, skin irritations, dry cough and sneezing (Mousavi et al., 2016; Mousavi et al., 2019). Several studies also reported a significant association between shortexposure to pollen and manifestations (Caillaud et al., 2014; Häfner et al., 2011; Kitinoja et al., 2020; Roberts et al., 2005). Therefore, the monitoring of airborne allergenic pollen grains can be helpful in predicting the severity of allergy seasons and appeared to be crucial for people suffering from pollinosis, estimated at 10-30% of the global population (Pawankar et al., 2011). By collecting and identifying pollen grains via appropriate methods and determining the time of their distribution in the air, a schedule could be applied and correlated to the date of allergy symptoms attack in sensitive people. This could help to determine the types of allergenic pollen grains occurring in each regions and to estimate the allergenic potential of different pollen species (Cariñanos González & Casares Porcel, 2019; Mansouritorghabeh et al., 2019).

Pollen calendars are used for graphic expression, timing, and pollen concentrations of different species in specific locations (Katotomichelakis et al., 2015). In Europe, long-term monitoring of pollen season of plant species is being carried out using a regular collection of relevant data, and many pollen monitoring stations in European countries are conducting regular and periodic surveys on airborne pollen grains (Sofiev et al., 2013). More than half of worldwide pollen monitoring stations are placed in this continent (598 out of 1013 active stations worldwide) (Buters et al., 2018). Most of them based on the Hirst pollen monitoring method (Hirst, 1952). The top European countries in terms of the number of active pollen monitoring stations are Italy (90 active stations), Spain (89 active stations), France (88 active stations) and Germany (58 active stations) (Buters et al., 2018); (https://www.zaumonline.de/pollen/pollen-monitoring-map-of-theworld.html, accessed12-08-2020). However, only a few countries in Europe, such as Switzerland, have government surveillance networks, and the majority of pollen and spore surveillance networks belong to the private sector, and in some cases their data are not freely available (Buters et al., 2018). In Asia, we can highlight the Japanese aerobiological network by its unique density (147 active stations). Although, historically the pollen time series are longer in other countries as India and Turkey. In Africa, most of the efforts to record airborne pollen are in the North African Mediterranean countries of Tunisia, Morocco and Algeria as well as in South Africa. In Oceania, a growing number of active stations are available in Australia since 2016 (21 active stations), following the dramatic thunderstorm event driven by airborne pollen causing 10 deaths (Thien et al., 2018). In America, pollen monitoring is performed routinely in most of the countries but the USA network shows the highest number (105 active stations) of recording stations (Fig. 1) (Buters et al., 2018; Hirst, 1952).

Pollen concentrations in an area are closely related to the distribution of plants in the area and are strongly influenced by climatic and meteorological factors (Lo et al., 2019). This review aimed at presenting relevant data that have been published on the influence of climate change and meteorological

factors on pollen season indicators (main pollen season, pollen production, pollen concentration, annual pollen integral, and pollination peak value) of main allergenic plant taxa. This article also provides insights on climate parameters influencing pollen production in main allergenic plant species and their possible impacts on human health. We performed a literature search of google scholar using the 150 key words and summarized more than 100 scientific articles related to pollen season indicators, meteorological parameters, and climate change. Given that the present article is not a systematic review, we may not have identified all studies, and we must acknowledge a certain publication bias.

2. Flowering phenology and pollen season indicators

Phenology is the timing of repetitive events in the life cycle of plants and animals in response to seasonal and climatic changes, therefore, it is one of the most important indicators of climate change (Schwartz, 2003). In relation to plants and health impacts, one of the most important phenological events is the emergence of the first flowers and the spread of pollen. Therefore, the concentration of pollen in the air is strongly regulated by phenology and flowering intensity, and data related to airborne pollen are widely used as phenological indicators (Aguilera et al., 2015; Cecchi et al., 2010; Chuine et al., 1998; García-Mozo et al., 2009; García-Mozo et al., 2016).




Figure. 1. Overview of the total number of active pollen monitoring stations worldwide. (Buters et al., 2018; https://www.zaum-online.de/pollen/pollen-monitoring-map-of-the-world.html, checked 12-08-2020).

The production and distribution of pollen grains are strongly regulated by internal and environmental factors such as regional climate, age and size of the plant, phenology and genotype. Thus, phenological studies represent an efficient tool to differentiate airborne pollen concentrations both temporally and spatially (Aguilera & Valenzuela, 2012; Branzi & Zanotti, 1992; García-Mozo et al., 2016; Scheifinger et al., 2013; Walther et al., 2002).

In aeropalynology, various indicators are used to describe the pollen season; mainly depending on their suspected use, being for aerobiologists or medical uses in the context of preventive programs (Pfaar et al., 2017). These indicators include main pollen season (duration of time when a specific pollen is present in the atmosphere in significant concentrations at a location), pollen production (quantity of pollen produced per anther in angiosperms), pollen concentrations (the number of airborne pollen grains per unit volume of air), annual pollen integral (the sum of daily pollen concentrations for a specific taxon over the pollen year), and pollination peak value (the maximum daily count recorded during a pollen season) (Galán et al., 2017; Ruiz-Valenzuela & Aguilera, 2018; Zhang et al., 2015). These pollen season indicators are the consequence of a combination of processes as pollen production, emission, and dispersion. Each one depends on a combination of plant endogenous factors, the environmental conditions, or the analyzed parameter. Annual pollen integral of allergenic pollen grains correlated with the severity of pollinosis symptoms among sensitized individuals and could be used to present inter-annual variations (Bastl et al., 2016).

3. The effect of climate change on flowering phenology and pollen season indicators

Climate, and especially the prevailing weather during the plant growth season, is the most important factor influencing plant phenology (Schwartz, 2003). The impacts of climate change on reproductive phenology and pollen season have been extensively studied by the analysis of longterm trends of pollen time series. Most of the analyses are based on simple linear regression between the parameters and the time, or decomposing the time series components (García-Mozo et al., 2014). So far, the short-term and longterm effects of environmental and meteorological factors such as light, temperature, rainfall, relative humidity, snow cover, hours of sunlight, and wind speed on the pollination period and concentration of airborne pollen grains have been well-documented. These effects vary depending on type of taxon and different climates

(D browska-Zapart et al., 2018; Donders et al., 2014; Emberlin et al., 2002; Galán et al., 2005). The pollen concentration is strongly influenced in hot and dry climates by average and maximum temperatures and water availability, while in semi-arid climates, it mainly relies on relative humidity, rainfall, hours of sunlight and daily temperature fluctuations (Rodriguez-Rajo et al., 2004). In the Mediterranean climate, pollen concentration is mainly impacted by rainfall and ambient temperature (Aboulaich et al., 2013; Uguz et al., 2017), in the semi-tropical climates by maximum temperature (Green et al., 2004) and in areas with temperate climates by relative humidity and maximum air temperatures (Puc, 2012).

Climate can affect the pollen production and pollination period, and consequently, the pollen concentration in the air. Several variations in the phenological behavior of plants over time, as a consequence of climate change have been reported. These changes are especially relevant for pollen exposure, especially those related to flowering phenology of anemophilous plants. For example, the International Phenological Gardens Network reported a six-day advance of the onset of spring events over a thirty-year period, with the highest rates of change in the Baltic region and Western Europe (Menzel & Fabian, 1999). It was found that climate change caused an up to six-day advance of phenological events in early spring and an up to five-day delay of autumn events in comparison with the early 1960s (Walther et al., 2002). Menzel (2000)reported a six-day advance for the flowering of some flowers over a 45-year period (1951-1996) in Europe (Menzel, 2000), and Fitter and Fitter (2002) have reported early flowering onset (about 4 to 5 days) for Great Britain over 10 years (Fitter & Fitter, 2002). In another study, Spieksma et al. (2003) analyzed several allergenic plant species to define their annual pollen integral in five cities located in Western Europe (Spieksma et al., 2003). Worryingly, data related to the highly allergenic birch pollen showed a rising trend in all studied stations and countries, although only two of the five rising trends were significant over time. A phenological study preformed in Switzerland between 1951 and 1998 showed a tendency towards earlier onset of the main pollen season in the spring and a weak tendency to start the main pollen season later in the fall (Defila & Clot, 2001). Bortenschlager and Bortenschlager (2005) reported that the flowering period of six taxa began earlier in Austria, thereby increasing the duration of the main pollen season of these species (Bortenschlager & Bortenschlager, 2005). In addition, pollination peak values and total pollen production showed an

increasing trend. In contrast, in Eastern Europe, phenological trends sometimes showed a delay of one to two weeks at the onset of spring events, mainly due to colder winters observed in these areas (Ahas et al., 2002).

Regarding pollen concentrations, an Italian study performed in Genova, found that the annual pollen concentration of *Parietaria* L. considerably increased from 1981 to 1997, while no significant increase in the pollen concentrations of *Artemisia* L. and Poaceae Barnhart were reported. (Voltolini et al., 2000). A longer duration of the pollen season has been also evidenced in Italy, peculiarly in summer (Beggs, 2004) as in the case of the Urticaceae Juss. (Frenguelli, 2002). In a grassy area in North America using simulated warming, rising temperatures led to the advance of main pollen season for species that flowered naturally before the peak of summer temperatures, and also delayed the

start of the main pollen season for species such as ragweed in which their flowering occurs after the peak of summer temperatures (Sherry et al., 2007). Based on a new method using dynamic systems, the effects of climate change on the concentration of airborne pollen grains can be predicted. Accordingly, de León et al. (de León et al., 2015) were able to predict an increase in the pollen concentration by 2070 from 28.5 to 44.3%, by examining the airborne pollen fluctuations of grass species from 1982 to 2012 in the city of Córdoba in southern Spain. The results of the studies investigating the effect of climate meteorological parameters on flowering phenology pollen season indicators of allergenic herbaceous taxa (grasses and weeds) are summarized in Table 1 and 2. Correlating data on aerobiological factors such as the onset, peak and main pollen season of allergenic species with pollen

Table 1. Studies investigating the effect of climate change and metrological factors on flowering phenology and pollen season indicators of allergenic herbaceous taxa (grasses).

Pollen Taxon	Parameter (s)	Location & time period	Phenology/ pollen season indicators	Reference (s)
Bromus japonicus, Dichanthelium oligosanthes, Panicum virgatum	Temperature	McClain (US) (2003-2004)	Main pollen season (Earlier flowering)	(Sherry et al., 2007)
Andropogon gerardii, Schizachyrium scoparium	Temperature	McClain (US) (2003- 2004)	Main pollen season (Delayed flowering)	(Sherry et al., 2007)
Poaceae	Temperature	Liguria (Italy) (1981- 2007)	Main pollen season (Increased duration, advanced onset date)	(Ariano et al., 2010)
Poaceae	North Atlantic Oscillation (NAO)	Iberian Peninsula (Spain and Portugal) (1994-2013)	Annual Pollen integral (Decreased)	(Galán et al., 2016)
Poaceae	Wind direction	Guadalajara (Spain) (2008-2013)	Pollen concentration (Concentration fluctuations)	(Rojo et al., 2015)
Poaceae	Temperature, Relative humidity, Wind speed, Radiation, Rainfall, Wind calm frequency	Brussels (Belgium) (1982-2015)	Pollen concentration (Decreased diurnally)	(Bruffaerts et al., 2018)
Poaceae	Temperature, Relative humidity, Wind speed, Radiation, Rainfall, Wind calm frequency	Brussels (Belgium) (1982-2015)	Main pollen season (Earlier flowering)	(Bruffaerts et al., 2018)
Poaceae	Temperature, Relative humidity	Šiauliai (Lithuania) (2010-2018)	Pollen concentration (Effected concentration)	(Sauliene et al., 2019)
Poaceae	Temperature, Solar irradiance, Sunlight	Guadalajara (Spain) (2008-2013)	Pollen concentration (Increased)	(Rojo et al., 2015)
Poaceae	Rainfall, Relative humidity	Guadalajara (Spain) (2008-2013)	Pollen concentration (Decreased)	(Rojo et al., 2015)

Table 2. Studies investigating the effect of climate change and metrological factors on flowering phenology and pollen season indicators of allergenic herbaceous taxa (weeds).

Pollen Taxon	Parameter (s)	Location & time period	Phenology/ pollen season indicators	Reference (s)
Ambrosia psilostachya	Temperature	Norman (Oklahoma) (1999-2001)	Pollen concentration (Increased)	(Wan et al., 2002)
Viola bicolor, Veronica arvensis, Cerastium glomeratum, Plantago virginica, Achillea millefolium, Erigeron strigosus	Temperature	McClain (US) (2003-2004)	Main pollen season (Earlier flowering)	(Sherry et al., 2007)
Ambrosia psilostachya	Temperature	McClain (US) (2003- 2004)	Main pollen season (Delayed flowering)	(Sherry et al., 2007)
Parietaria	Temperature	Liguria (Italy) (1981-2007)	Main pollen season (Increased duration, advanced onset date) Pollen concentration (Increased)	(Ariano et al., 2010)
Plantago major, Urticaceae	Temperature	Jaen (Spain) (1994-2016)	Pollen concentration (Decreased)	(Ruiz-Valenzuela & Aguilera, 2018)
Artemisia	Temperature	Poznan (Poland) (1995-2004)	Pollen concentration (Increased daily)	(Stach et al., 2007)
Artemisia	Relative humidity	Poznan (Poland) (1995-2004)	Pollen concentration (Decreased daily)	(Stach et al., 2007)
Ambrosia	Latitude	Maryland (US) (1995-2009)	Main pollen season (Increased duration)	(Ziska et al., 2003)
Artemisia	Rainfall	Córdoba (Spain) (1991-2011)	Main pollen season (Earlier flowering, Increased duration)	(Cariñanos et al., 2014)
Amaranthaceae	Rainfall	Iberian Peninsula (Spain and Portugal) (1994-2013)	Annual pollen integral (Increased)	(Galán et al., 2016)
Urticaeae, <i>Plantago</i> , <i>Rumex</i> , Amaranthaceae, <i>Artemisia</i>	North Atlantic Oscillation (NAO)	Iberian Peninsula (Spain and Portugal) (1994-2013)	Annual pollen integral (Decreased)	(Galán et al., 2016)
Chenopodiaceae– Amaranthaceae, Urticaceae, <i>Plantago</i> , <i>Rumex</i>	Wind direction	Guadalajara (Spain) (2008-2013)	Pollen concentration (Concentration fluctuations)	(Rojo et al., 2015)
Artemisia vulgaris	Temperature, Relative humidity, Wind speed, Radiation, Rainfall, Wind calm frequency	Brussels (Belgium) (1982-2015)	Pollen concentration (Decreased diurnally)	(Bruffaerts et al., 2018)
Urticaceae	Temperature, Relative humidity, Wind speed, Radiation, Rainfall, Wind calm frequency	Brussels (Belgium) (1982-2015)	Main pollen season (Earlier flowering)	(Bruffaerts et al., 2018)
Artemisia	Temperature, Relative humidity	Šiauliai (Lithuania) (2010-2018)	Pollen concentration (Effected concentration)	(Sauliene et al., 2019)
Chenopodiaceae– Amaranthaceae, Urticaceae, <i>Plantago</i> , <i>Rumex</i>	Temperature, Solar irradiance, Sunlight	Guadalajara (Spain) (2008-2013)	Pollen concentration (Increased)	(Rojo et al., 2015)
Chenopodiaceae– Amaranthaceae, Urticaceae, <i>Plantago</i> , <i>Rumex</i>	Rainfall, Relative humidity	Guadalajara (Spain) (2008-2013)	Pollen concentration (Decreased)	(Rojo et al., 2015)

concentration data can be used as evidence for changes in the flowering and onsets of main pollen season as well as for the prediction of pollen season of the species of interest (Clot, 2003; Emberlin et al., 2002; Galán et al., 2005; Huynen, 2003; Rasmussen, 2002; Van Vliet et al., 2002). In this regard, the majority of studies have focused on allergenic plants such as Betula L. (Emberlin et al., 2002; Van Vliet et al., 2002), Artemisia vulgaris L. (Stach et al., 2007), Urticaceae Juss. (Frenguelli, 2002), grasses (Burr, 1999; Emberlin et al., 1999), Quercus L. (García-Mozo et al., 2006; Van Vliet et al., 2002), Cedrus Mill. (Tosunoglu et al., 2015) and Platanus L. (Tedeschini et al., 2006). Some studies have predicted the advance of the onset of the main pollen season from one to three weeks for Olea europea L. (Galán et al., 2005) and one month for oak by the end of the century (García-Mozo et al., 2006). Also, the increase in spring temperature in the Mediterranean region leads to earlier flowering of olive trees (Garcia-Mozo et al., 2009; Orlandi et al., 2010). Likewise, in Italy, prediction models point towards a prolonged pollination period for olive trees throughout the 21st century (Avolio et al., 2012). Spieksma et al. (1995) examined the birch pollen concentration in the European atmosphere and reported quantitative annual fluctuations and changes in onset dates of the main pollen season (Spieksma et al., 1995). This study revealed that the main pollen season of the birch tree shows a tendency towards an earlier onset, which has nothing to do with the observed weather trends. Likewise, Emberlin et al.'s research (1997) on the onset of the main pollen season of birch in three regions of the United Kingdom over a 42-year period showed a tendency for the main pollen season to onset earlier (Emberlin et al., 1997). However, in this latter study, this trend was closely related to higher temperature records (more than 5.5 degrees centigrade) from January to March with 3month cumulative temperatures ranging from 25 to 30 degrees centigrade for the study sites. Interestingly, the main pollen seasons of birch have advanced in the studied areas by about Five days per decade. In another study, Emberlin et al. (2002) investigated over a 20-year period the correlation between the temporal patterns of change in spring temperatures and changes in the onset of the main pollen season of birch in several selected areas in Europe (Brussels, Kevo, London, Turku, Vienna and Zurich), in order to predict these patterns for the next ten years. Accordingly, the city of Kevo showed a six-day delay, the cities of London, Brussels, Zurich, and Vienna showed about a sixday advance, and the city of Turku showed a cyclical pattern (Emberlin et al., 2002). Early

flowering of *Bromus rubens* Delile. and *Hordeum leporinum* Link., that routinely bloom in in start of spring, in years with higher mean winter temperatures and *Trisetaria panicea* (Lam.) Paunero and *Dactylis glomerata* subsp. *hispanica* (Roth) Nyman, that routinely bloom in mid to late spring in years with higher cumulative rainfall in spring and winter has been reported in city of Toledo, Spain (Romero-Morte et al., 2020).

Some studies have examined simultaneously a wide range of pollen flora in a specific area. For example, Clot's study showed that 71% of the main pollen season onset or end dates show a significant advance, and for the majority of pollen species, the main pollen season does not last longer but changes over time(Clot, 2003). Both Damialis et al. (2007) and Cristofori et al. (2010) reported a trend towards a significant increase of pollen concentrations for the majority of plant taxa. In these studies, tree species showed a significant increase in their pollen production compared to herbaceous plants (Damialis et al., 2007) (Cristofori et al., 2010) . Table 3 provides a summary of studies investigating the effect of climate and meteorological parameters on flowering phenology and pollen season indicators of allergenic tree taxa.

Investigation of the main pollen season changes in allergenic species of birch (*Betula*), oak (*Quercus*), mugwort (*Artemisia*), and grass (Poaceae) using registered data related to daily airborne pollen grains and climatic factors from 1994 to 2010 across the contiguous USA has shown that the main pollen season of trees, grasses, and weeds during the 2001-2010 decade has advanced three days on average when compared to the previous decade (1994-2000). Also, the average peak value and the total number of annual airborne pollen recorded daily were 42.2% and 46% higher, respectively (Zhang et al., 2015).

During an 18-year study, Scevkova et al. (2021) examined the duration and intensity of the pollen season in three allergenic taxa (*Alnus* Mill., Poaceae, *Artemisia*). During this period, the flowering period was shorter for Poaceae and *Artemisia*. The pollination peak reached earlier for *Alnus* and *Artemisia*. *Alnus* peak value showed a significant rising trend and the duration of pollen season and peak values of *Artemisia* and Poaceae had a declining trend (Š evková et al., 2021).

Seasonal variations of airborne pollen concentrations can be related to phenological and/or meteorological factors. In this regards, a quantitative evaluation of airborne pollen in during 2018 to 2020 showed that the highest concentration of airborne pollen is related to spring and autumn. This could be due to the favorable phenological and meteorological factors for plant growth, dispersion and pollen transfer in these seasons (Ravindra et al., 2021).

Table 3. Studies investigating the effect of climate change and metrological factors on flowering phenology and pollen season indicators of allergenic arboreal taxa (trees).

Pollen Taxon	Parameter (s)	Location & time period	Phenology/ pollen season indicators	Reference (s)
Betula	Temperature	Basel (Switzerland) (1969-2006)	Main pollen season (Earlier flowering) Annual pollen integral (Increased) Pollination peak value (Increased)	(Frei & Gassner, 2008)
Betula, Cedrus, Olea	Temperature	Liguria (Italy) (1981-2007)	Main pollen season (Increased duration, advanced onset date) Pollen concentration (Increased)	(Ariano et al., 2010)
Betula	Temperature	(Finland) (1974-2004)	Pollen concentration (Increased)	(Yli-Panula et al., 2009)
Cupressaceae, Olea, Pinus, Platanus, Quercus	Temperature	Jaen (Spain) (1994-2016)	Main pollen season (Increased duration) Pollen concentration (Increased)	(Ruiz-Valenzuela & Aguilera, 2018)
Betula	Altitude	Zugspitze (Germany) (2009-2010)	Pollen concentration (Decreased)	(Jochner et al., 2015)
Pinus taeda	Carbon Dioxide	North Carolina (US) (1996-2004)	Main pollen season (Advanced onset date, prolonged pollen season) Pollen concentration (Increased)	(LaDeau & Clark, 2006)
Cupressaceae, Quercus, Platanus, Olea, Populus, Pinus, Ulmus, Moraceae, Fraxinus	Wind direction	Guadalajara (Spain) (2008-2013)	Pollen concentration (Concentration fluctuations)	(Rojo et al., 2015)
Alnus, Corylus, Betula, Carpinus, Quercus, Fraxinus, Platanus	Temperature, Relative humidity, Wind speed, Radiation, Rainfall, Wind calm frequency	Brussels (Belgium) (1982-2015)	Pollen concentration (Increased diurnally)	(Bruffaerts et al., 2018)
Betula, Quercus, Fraxinus, Platanus	Temperature, Relative humidity, Wind speed, Radiation, Rainfall, Wind calm frequency	Brussels (Belgium) (1982-2015)	Main pollen season (Earlier flowering)	(Bruffaerts et al., 2018)
Quercus	Temperature, Radiation, Wind speed, Wind frequency	Malaga (Spain) (1992-2015)	Pollen concentration (Increased diurnally and weekly)	(Recio et al., 2018)
Quercus	Rainfall, Relative humidity	Malaga (Spain) (1992-2015)	Pollen concentration (Decreased diurnally and weekly)	(Recio et al., 2018)
Alnus, Betula, Corylus	Temperature, Relative humidity	Šiauliai (Lithuania) (2010-2018)	Pollen concentration (Effected concentration)	(Sauliene et al., 2019)
Cupressaceae, Quercus, Platanus, Olea, Populus, Pinus, Ulmus, Moraceae, Fraxinus	Temperature, Solar irradiance, Sunlight	Guadalajara (Spain) (2008-2013)	Pollen concentration (Increased)	(Rojo et al., 2015)
Cupressaceae, Quercus, Platanus, Olea, Populus, Pinus, Ulmus, Moraceae, Fraxinus	Rainfall, Relative humidity	Guadalajara (Spain) (2008-2013)	Pollen concentration (Decreased)	(Rojo et al., 2015)

3.1 Effect of temperature

Flowering phenology in many plant species appeared to be closely related to temperature when compared to other metrological parameters (Moore & Lauenroth, 2017). The impact of temperature on plants reproductive behavior depends on the

exposure time along the reproductive cycle (Rojo et al., 2020). Also, it is essential to take into account that the impact of weather conditions depends on the state of the plants and the range of values (Oteros et al., 2013a; Oteros et al., 2013b). In tree species, flowering phenology is specifically regulated by

temperature and in herbaceous species, it is regulated by temperature and water availability (García-Mozo et al., 2016). Flowering phenology of plants pollinating in spring and early summer is strongly influenced by temperature, while the photoperiod was found to be more important for plants flowering from late summer to autumn (Beaubien & Freeland, 2000). In addition to the variation of temperatures in spring and increased global temperatures leading to milder winters and warmer springs may result to the early flowering of many spring species in different regions (Fitter & Fitter, 2002; Linderholm, 2006; Menzel, 2000). Based on the Marsham family phenological data record (1736), Margary suggested that if spring average temperatures increased by one degree centigrade, a 5- to 7-day advance in spring events would be expected (Menzel, 2000).

Studies on ragweed (*Ambrosia psilostachya* DC.) showed that, when plants are exposed to high temperatures, the number of their stems increased by 88%, which increases the pollen production rate in each pot by 84% (Wan et al., 2002). In another study, Sherry et al. (2007) looked at the effects of rising ambient temperatures on the flowering of a number of species and reported early flowering in species that flower in spring and early summer (Sherry et al., 2007). However, species that flowered naturally in late summer and autumn showed a delay in flowering.

Frei and Gassner (2008) used data from long-term airborne pollen data in Basel, Switzerland, for a period of 38 years and reported that flowering season in birch occurs 15 days earlier due to the increase in the temperature and also, there was a tendency to increase the annual pollen integral and the daily peak value (Frei & Gassner, 2008).

Ariano et al., in a 27-year long-term study (between 1981 and 2007), studied the effect of climate changes, especially rising temperatures, on the pollen season of 5 major allergenic pollen (birch, olive, cedar, *Parietaria* and grass) in Liguria (Italy) (Ariano et al., 2010). The results showed that duration of the main pollen season increased for Parietaria spp. (85 days), cedar and olive (18 days each). In all mentioned species, a general advance of the main pollen season was also observed. The pollen concentrations of all intended species other than grasses also increased by an average of about 25 percent. These behaviors were parallel to the increase in the continuity of direct radiation, temperature, and the number of days with temperatures above 30 degrees centigrade. A longterm study performed by Yli-Panula et al. (2009) demonstrated that high temperatures promote early phenological advance and increase concentrations of

birch airborne pollen during a 31-year period in the Turku region of Finland. However, no changes were reported in the main pollen season (Yli-Panula et al., 2009).

According to the Intergovernmental Panel on Climate Change, the increase in temperatures is closely associated with the latitude. Ziska et al. (2003) investigated and compared the main pollen season of the ragweed from 1995 to 2009 in North America. Their data showed a significant increase in the main pollen season of ragweed from 13 to 27 days at latitude above 44 ° N (Ziska et al., 2003).

A study investigating changes in the pollen spectrum of twelve types of allergenic airborne pollens from southern Spain over a 23-year period by Ruiz-Valenzuela and Aguilera (2018) that, with increasing temperature, the pollen concentration of species increases, while the pollen concentrations in grass and weed taxon's especially Plantago major L., shows a decreasing trend (Ruiz-Valenzuela & Aguilera, 2018). Therefore, the main pollen season of tree species appears to be more affected by temperature fluctuations and showed significant changes in response to climate change. The main pollen season of these tree species also shows a tendency to become longer. The results of several researches suggested that Quercus, Platanus, Pinus L. and Cupressaceae Gray pollen grains, along with Plantago L. pollen grains could be used as efficient biological indicators for identifying local climate change (Ziska et al., 2019). In various geographic locations across the northern hemisphere, recent investigations most demonstrated that an increase in temperature (minimum and maximum temperatures) significantly contributed to increased pollen concentrations and extended the main pollen season for numerous airborne allergic pollen species (Ziska et al., 2019). Some studies also point other factors closely related to temperature, like slope orientation or altitude, as relevant driver of flowering phenology e.g. (Oteros et al., 2013c)

3.2. The Effect of carbon dioxide

High levels of CO2 may increase the concentration of airborne pollen grains in large cities and thus increase respiratory allergies. Carbon dioxide from human activities, as a major factor implicated in climate change, has direct (stimulating photosynthesis and plant growth) and indirect (increasing the average surface temperature of the Earth) effects on plants (Ziska et al., 2012). Pioneer researches showed an increase in the atmospheric concentrations of carbon dioxide, about 200 ppm above the ambient atmosphere values (600 ppm), highly stimulated photosynthesis by up to 60% in

some C3 plants (Curtis & Wang, 1998; Jablonski et al., 2002; Norby et al., 1999). This change in carbon adsorption may directly or indirectly impact the physiology, phenology the reproductive behavior and the geographical distribution of plant species. As a result, the plant's biomass, the number of flowers and pollen production significantly increased, and subsequently, the concentration of allergenic airborne pollen is affected (Jablonski et al., 2002). For example, some studies performed on the invasive and highly allergenic ragweed plant have reported a significant increase in the number of flower spikes per bush and a 60 to 90 percent increase in pollen production under an approximate doubling of carbon dioxide concentrations (Rogers et al., 2006; Singer et al., 2005; Wayne et al., 2002). A study on one of the fastest-growing pine species (Pinus taeda L.), extensively planted worldwide has shown that increasing concentrations of carbon dioxide led to an elevated pollen production as well as to the early onset and prolongation of the main pollen season (LaDeau & Clark, 2006). Likewise, rising temperatures and carbon dioxide appeared to significantly increase pollen production in many common species (Blando et al., 2012).

Other studies showed that an enrichment with 500 to 800 ppm carbon dioxide of plant reproductive organs led to a 31% increase of the total plant mass and the intensity of flowering was 19% higher in treated crop and wild plant species in comparison with controls (Curtis & Wang, 1998; Jablonski et al., 2002). In one study, Knapp and Soulé (1998) studied the spread of Juniperus occidentalis Hook. populations under increased amounts of atmospheric carbon dioxide over a 23-year period in Oregon (Knapp & Soulé, 1998). Their results showed that increasing atmospheric carbon dioxide would preferably increase the growth and spread of young trees. In addition, similar studies have suggested that ragweed quantitative and qualitative spread during the twentieth century has been due to increased concentrations of atmospheric carbon dioxide (Singer et al., 2005; Ziska & Caulfield, 2000).

3.3. The effect of rainfall

Rainfall can have a dual effect on the concentration of airborne pollen. Like temperature, the impact of rainfall depends directly on the exposure time along the reproductive cycle. In some studies, the rain during the main pollen season has reduced the pollen concentration in the atmosphere, thereby reducing the allergy symptoms caused by herbaceous pollen grains (Ursu, 2012). In contrast, after heavy rains, increased concentrations of pollen, other bio-aerosols and free allergen molecules binding airborne particles have been reported

(Huffman et al., 2013; Müller-Germann et al., 2015; Schäppi et al., 1997; Taylor et al., 2007). It appeared that in a warmer climate, heavy rains will increase and tend to fewer but more intense events. In this respect, it comes that heavy rainfalls and strong winds increased asthma attacks among pollen allergic individuals, a phenomenon repeatedly reported and known as thunderstorm asthma in many regions worldwide (Girgis et al., 2000; Suárez-Varela et al., 2008).

In Poland, the flowering phenology of *Artemisia* pollen as one of the most important causes of respiratory allergies appeared to be affected by climate change, especially rainfall. In a study on the concentration of *Artemisia* airborne pollen grains between 1995 and 2004 in southern Poland (Poznan), Stach et al. (2007) reported that the main pollen season of *Artemisia* is greatly affected by rainfall in weeks preceding the pollen season, which the main pollen season starts earlier and lasts longer. The authors also reported a direct effect of the temperature on the daily pollen concentration in this genus, while relative humidity has the opposite effect on its pollen concentration (Stach et al., 2007).

Cariñanos et al. (2014) investigated a 21-year data set (1991-2011) related to airborne pollen grains of Amaranthaceae Juss. in Córdoba, in the southern Iberian Peninsula, to assess the impact of climatic conditions on pollen season indicators. This study revealed the very long persistence (from early spring to early autumn) of Amaranthaceae pollen in the atmosphere. The annual pollen integral showed a significant increase (from approximately 200 pollen grains to about 2,000 pollen grains) which was strongly affected by rainfall during the flowering period, causing the growth of new plants and thus increased pollen production (Cariñanos et al., 2014).

In southern Europe, lack of access to water may lead to a decline in flowering, especially in herbaceous species (Alcázar et al., 2009; Recio et al., 2009). As North Atlantic Oscillation (NAO) also affects rainfall (Gallego et al., 2005), Galán et al. (2016) examined over two decades (1994-2013) the effect of NAO on the flowering intensity of 12 inflated taxa at 12 locations of the Iberian Peninsula by analyzing and comparing their annual pollen index (API). Their results showed a negative relationship between the pollen concentration in the atmosphere and the NAO index averages during winter(Galán et al., 2016). These results suggested that the phenology of many Mediterranean species is influenced by variations in rainfall, partly driven by climate change.

3.4. The effect of atmospheric pollutants

Among the factors that may affect the concentration of airborne pollen grains are atmospheric pollutants in urban areas such as suspended particles in the atmosphere (PM, coarse and fine), ozone (O3), carbon monoxide (CO), dioxide nitrogen (NO2) and sulfur dioxide (SO2) (Sauliene et al., 2019). Orby et al. (2015) showed that there was a significant association between pollen concentrations and atmospheric ozone (Ørby et al., 2015). Sauliene et al. (Sauliene et al., 2019) also showed that the concentrations of O3 and PM10 pollutants were significantly associated with the concentration of airborne pollen grains during the main pollen season. The SO2 pollutant showed a positive correlation with Betulaceae Grav pollen concentrations and a negative correlation with grasses pollen concentrations. Recently, Oduber et al. (2019) studied the effect of air pollutants concentration on airborne pollen grains concentration. The researchers looked at the association of pollen concentration trends of allergenic species i.e Fraxinus Tourn. ex L., Poaceae, and Populus L. with the concentration trends of O3, NO2, NO, CO, SO2 and, PM10 atmospheric pollutants over the past two decades in the city of Leon. In this study, a significant decreasing trend was observed in the concentration of air pollutants, while the concentration of Fraxinus airborne pollens a significantly decreased. No tendency was reported for the advance or delay of the main pollen season of any of the three taxa. Analysis of their data using the Spearman correlation method showed that the duration of flowering and pollination is highly dependent on the weather conditions before these periods and is affected by the concentrations of atmospheric pollutants. However, they were unable to find any significant relationship statistically between airborne pollen concentrations and air pollution during pollination periods (Oduber et al., 2019). Therefore, although the effect of air pollutants on the allergen content of pollen grains is well documented (Chassard et al., 2015; Frank & Ernst, 2016; Mousavi et al., 2019; Sénéchal et al., 2015 (Shahali et al., 2009), it does not appear to be a clear relationship between air pollution parameters and the concentration of airborne pollutants, although further research is needed.

3.5. The effect of land use

One of the important reasons for the difference in flowering phenology and pollen season indicators among plant species and in different parts of the world is the difference in land use (Scheifinger et al., 2013). Land use determines the number of individuals of a plant species in a given area

(García-Mozo et al., 2016). As the airborne pollen grains identified in a given area is directly related to local land uses, this factor can strongly influence changes in the pollen spectrum and concentration of airborne pollen grains (Galán et al., 2016). García-Mozo et al. (2016) investigated the common influence of climate variables and land use change on airborne pollen concentrations fluctuations in the city of Cordoba (southern Spain) over a 15-year period. A significant increase in some pollen types such as olives and grasses was found, while the airborne pollen concentration of some invasive taxa such as Amaranthaceae, Rumex L., Plantago, Urticaceae decreased, which could be related to fluctuations in vegetation, climate or land use change (García-Mozo et al., 2006). In another study, using comprehensive modeling methods, Hamaoui-Laguel et al. (2015) estimated that by 2050, the concentration of ragweed airborne pollen grains, an allergenic invasive species in Europe (Kazinczi et al., 2008), would be four times higher (Hamaoui-Laguel et al., 2015). This prediction largely depends on the hypotheses of the pollen distribution rate. About two-thirds of the ragweed airborne pollen concentration is related to land use change and climate change. Therefore, land use change and climate change could lead to the expansion of ragweed habitats in northern and eastern Europe, thereby increasing ragweed pollen production levels in these areas in the coming years (Kazinczi et al., 2008).

3.6. The effect of urbanization

The physical properties of the lower layers of the atmosphere could be modified both horizontally and vertically by urban environments. This leads to a change in the equilibrium of solar radiation and creates an urban heat island which is the term used to describe the dome of warm air covering large cities (Cecchi et al., 2010). This phenomenon, which is characterized by higher temperatures, lower air humidity levels, and warm winds t, causes pollen deposition and changes in pollen distribution patterns in cities and prolonged plant growing periods and thus increases airborne pollen grains concentration (Mimet et al., 2009; Unger, 1999). This phenomenon leads to the pollen season advance in urban areas compared to rural areas in about 2 to 4 days (D'Amato, 2011). Also, in cities, the increase in turbulence caused by specific thermal wind patterns can transfer pollen grains from suburban areas to urban centers (Emberlin & Norris-Hill, 1991). In urban areas, earlier flowering and pollination have been reported for Ambrosia L. compared to rural areas (Ziska et al., 2003). The same observations have been reported for some

allergenic tree species such as Platanus acerifolia (Aiton) Willd. (Emberlin & Norris-Hill, 1991) and a number of non-allergenic plants. However, in the case of Ambrosia spp., rising temperatures and high concentrations of carbon dioxide in urban areas have led to a significant reduction in the urban population of this invasive species (Ziska et al., 2007). There is evidence that urbanization can significantly reduce herbaceous pollen grains such as grasses (Cecchi et al., 2010). The declining trend in the number of grass annual pollen related to urbanization in Leiden (Netherlands) (Jäger et al., 1991), London and Derby (UK) (Emberlin et al., 1999; Frenguelli, 2002), Parma (Italy) (Ridolo et al., 2007), and even lower pollen peaks and fewer pollination days have been reported in other studies (Jato et al., 2009; Minero et al., 1998).

3.7. Combined effect of several parameters

Some studies have studied the simultaneous effect of several meteorological parameters on the concentration of airborne pollen grains. example, a study performed between 1982 and 2015 in Brussels (Bruffaerts et al., 2018) examined the relationship between pollen concentrations and meteorological parameters. Using daily obtained from 11 pollen types (8 tree species and 3 herbaceous species) and 10 meteorological parameters, a general increasing trend in daily airborne pollen grains of tree species (except for European beech species) and a general decreasing trend in daily airborne pollen grains for herbaceous species (except Urticaceae) were reported. Early flowering was also observed for birch, oak, ash, plane tree, grass, and Urticaceae trees. The rate of change in the annual cycle of several meteorological parameters such as rainfall, humidity temperature and radiation, was significantly associated with the rate of change in the annual cycle of airborne pollen grains (Bruffaerts et al., 2018).

In oak, the formation of flower buds is very sensitive to temperature. In this regard, a 25-year aerobiological study of the effect of climate variables on oak pollen atmospheric concentrations in the coastal city of Malaga (southwestern Mediterranean basin) showed that fluctuations in oak pollen atmospheric concentrations (daily and weekly values) are significant and has a positive relationship with temperature, changes in solar radiation, wind speed, and northwest wind frequency, and a negative relationship with precipitation and relative humidity. The peak of oak pollen production was observed almost every four years, which corresponds to periods of drought. These results suggest that the reason for the observed trend in oak pollen production is probably

the increase in atmospheric temperature and drought, which is increasing in the western Mediterranean region (Recio et al., 2018). Sauliene et al. (2019) examined the relationship between the airborne pollen concentration of three taxa of the birch genus (Alnus, Betula and Corylus L.), grasses, and Artemisia genus with a set of meteorological factors and air pollution levels in Lithuania between 2010 and 2018. The results showed that the relative humidity and air temperature parameters had a significant effect on the airborne pollen concentrations of Alnus, Artemisia, Betula, Corylus and Poaceae. However, wind speed has little to do with the concentration of these airborne pollutants. Temperature conditions had the greatest impact on the concentration of *Corylus* and *Alnus* pollen grains (both early in flowering). There was no statistically significant relationship between Corylus pollen concentrations and air pollution. Betula and grasses pollen concentrations also had a positive correlation with ozone pollution (Sauliene et al., 2019).

Wind speed and direction (Damialis et al., 2005; Palacios et al., 2000) is one of the most important determinants of the distribution and transmission of airborne pollen grains. Small pollen grains are able to reach the upper layers of the atmosphere and travel long distances (Hernández-Ceballos et al., 2011; Rojo & Pérez-Badia, 2015; Smith et al., 2008). Most pollen grains do not travel a long distance from the emission area and are often distributed locally (Lavee & Datt, 1978). Therefore, the pollen spectrum provides important indications regarding the vegetation in/close to the area (Hamaoui-Laguel et al., 2015; González Minero & Candau, 1997). Rojo et al. (Rojo et al., 2015) examined and analyzed the pollen spectrum of the city of Guadalajara (central Spain) up to a radius of 20 km from the perspective of vegetation and land use in order to identify the source of pollen in the air further explore the potential effect of meteorological parameters on pollen propagation. This study revealed that the local wind direction was one of the most important variables influencing the concentration of different types of airborne pollen grains, and land use played a key role in airborne pollen counts and distribution in urban green spaces. The propagation of pollen grains from ornamental species was strongly influenced by eastern winds (plane tree), southern winds (cedar), western winds (cedar and plane tree) of the areas where the largest parks and gardens of the city were located. Cedar pollen is basically transmitted by the wind from the eastern edge of the city. Most pollen counts of Populus were related to western and eastern winds in environments containing rivers and streams. Temperatures, hours of solar irradiance and sunlight had a positive effect on the number of pollen grains in the air, while rainfall and relative humidity had a negative effect.

A study performed by Alan et al. (2018) tried to determine over a two-year period the relationship between airborne concentrations of grass pollen grains and Phl p 5 concentrations (group 5 of grass pollen allergens) in two cities with different climatic and geographical characteristics (Ankara and Zongledok). It was found that the index of the total number of grass pollen and the total concentration of Phl p 5 allergen in these two regions was different. The temperature appeared to be the main meteorological factor that affects the airborne pollen concentration and the Phl p 5 allergen concentration at both stations. Rainfall was also important for Zonguldak because of its geographical and climatic characteristics. Finally, the researchers suggested the term "drift effect" to describe the critical impact of the initial wind direction on the propagation of airborne allergens. Therefore, the distribution of allergen and pollen grains is highly influenced by wind direction, especially in areas with elevated topography (Alan et al., 2018). In this regard, Jochner et al. (2015) showed that pollen and allergen concentrations were significantly lower at higher altitudes (Jochner et al., 2015).

Conclusion

Phenological monitoring is a useful tool for assessing the effects of climate change on plants behavior. In recent decades, climate change has led to the advance of spring events and the delay of autumn phenological events. Considering reproductive phenology, most of the impacts are related to the advance of the flowering. This can affect the pollen season indicators of the species and, consequently, the airborne pollen concentrations and risk of pollen exposure. Flowering phenology and pollen season indicators vary for allergic plant species according to their geographical location and is strongly influenced and meteorological climatic parameters. Temperature, wind speed and direction, humidity, and rainfall are among the most important meteorological parameters affecting the fluctuations of annual concentrations of airborne pollens. Interestingly, tree species appeared to be particularly sensitive to temperature fluctuations and temperature is the most important climatic parameter affecting the flowering phenology and pollen season indicators of anemophilous trees releasing abundant concentrations of allergenic pollen grains. The most important pollen season indicators that are affected by temperature include annual pollen integral, main pollen season and pollen production (Fig. 2). Flowering phenology

of plant species that flower in spring and early summer is strongly affected by temperature, while species that flower in late summer and fall are generally more significantly impacted by photoperiod or lighting. Rainfall can affect some indicators of the pollen season, such as annual pollen integral, main pollen season and pollen concentration. However, elevated CO2 has an enhancing impact on pollen production, pollen concentration and main pollen season (Fig. 2).

The change in the main pollen season and airborne pollen concentration also depends on latitude, which is directly linked to temperature changes and rainfall during plant growth. For allergenic tree, grass and weed species, the main pollen season can be predicted by combining long-term data of pollen concentrations with onset time and pollination peak value. It is now well-documented that increasing global temperature due to climate change could lead to warmer winters and springs, resulting in the early flowering of many plant species in different parts of the world. There is also a growing body of scientific evidence supporting the fact that carbon dioxide from human activities results in increased plant biomass, increased flowering intensity, and pollen production by stimulating photosynthesis and plant growth and through its influence on the average temperature of the Earth's surface (Fig. 2). However, so far, no statistically significant relationship has been observed between the concentration of airborne pollen grains and the concentration of industrial/ urban air pollutants and further studies are still required to draw any conclusion regarding the exceedingly complex interactions between air pollution and the concentration of allergic pollen grains in different areas. The present review provides a global view of the effects of climate change and meteorological factors on flowering phenology and pollen season indicators of allergenic plant taxa worldwide, paving the way for comprehensive studies in this area of major environmental and public health importance.

Acknowledgements

The authors would like to thank Matthias Werchan from Charité Universitätsmedizin Berlin for his comments on the early version of this article. We also thank Prof. Lewis Ziska from USDA's Agricultural Research Service (Washington, D.C., United States) for Critical review of the article. This study was supported by the grant [ARI-99-20-PGA-IP-1-1] from space Biology and Environment center, Aerospace Research Institute (ARI), Ministry of Science Research and Technology (MSRT), Tehran, Iran.




Figure. 2. A Schematic outline of climate change and meteorological factors affecting pollen season indicators and pollen exposure. The long term and short-term effects of temperature, carbon dioxide, and rainfall on flowering phenology, pollen season indicator and pollen exposure are depicted. Pollen season indicators that have been studied include pollen production (quantity of pollen produced per anther in angiosperms), pollen concentration (the number of airborne pollen grains per unit volume of air), main pollen season (the presence of a specific pollen in the atmosphere in significant concentrations at a location) and annual pollen integral (the daily pollen concentration for a specific taxon over the pollen year). Increasing these indices, especially annual pollen integral, can increase the risk of pollen exposure in sensitized individuals. The increase in atmospheric carbon dioxide concentrations emitted through anthropogenic activities can amplify greenhouse effects, which elevate the Earth's surface temperature (global warming). This abnormal rise in global temperature could lead to climate change and increased heavy precipitation events. The up and down arrows indicate enhancing and lowering impacts, respectively and dashed arrows shows the potential impact.

REFERENCES

Aboulaich, N., Achmakh, L., Bouziane, H., Trigo, M.M., Recio, M., Kadiri, M., Cabezudo, B., Riadi, H. & Kazzaz, M. 2013. Effect of meteorological parameters on Poaceae pollen in the atmosphere of Tetouan (NW Morocco). International Journal of Biometeorology 57: 197-205.

Aguilera, F., Orlandi, F., Ruiz-Valenzuela, L., Msallem, M. & Fornaciari, M. 2015. Analysis and interpretation of long temporal trends in cumulative temperatures and olive reproductive features using a seasonal trend decomposition procedure. Agricultural and Forest Meteorology 203: 208-216.

Aguilera, F. & Valenzuela, L.R. 2012. Microclimatic-induced fluctuations in the flower and pollen production rate of olive trees (*Olea europaea* L.). Grana 51: 228-239.

Ahas, R., Aasa, A., Menzel, A., Fedotova, V. & Scheifinger, H. 2002. Changes in European spring phenology. International Journal of Climatology: A Journal of the Royal Meteorological Society 22: 1727-1738.

Alan, ., ahin, A.A., Sarı ahin, T., ahin, S., Kaplan,
A. & Pınar, N.M. 2018. The effect of geographical and climatic properties on grass pollen and Phl p 5

allergen release. International Journal of Biometeorology 62: 1325-1337.

Alcázar, P., Stach, A., Nowak, M. & Galán, C. 2009. Comparison of airborne herb pollen types in Córdoba (Southwestern Spain) and Poznan (Western Poland). Aerobiologia 25: 55-63.

Ariano, R., Canonica, G.W. & Passalacqua, G. 2010. Possible role of climate changes in variations in pollen seasons and allergic sensitizations during 27 years. Annals of Allergy, Asthma & Immunology 104: 215-222.

Avolio, E., Orlandi, F., Bellecci, C., Fornaciari, M. & Federico, S. 2012. Assessment of the impact of climate change on the olive flowering in Calabria (southern Italy). Theoretical and Applied Climatology 107: 531-540.

Bastl, K., Kmenta, M., Pessi, A.M., Prank, M., Saarto,
A., Sofiev, M., Bergmann, K.C., Buters, J. T.,
Thibaudon, M. & Jäger, S. 2016. First comparison of symptom data with allergen content (Bet v 1 and Phl p 5 measurements) and pollen data from four European regions during 2009–2011. Science of the Total Environment 548: 229-235.

Beaubien, E. & Freeland, H. 2000. Spring phenology trends in Alberta, Canada: links to ocean temperature.

- International Journal of Biometeorology 44: 53-59.
- **Beggs, P.J.** 2004. Impacts of climate change on aeroallergens: past and future. Clinical & Experimental Allergy 34: 1507-1513.
- **Blando, J., Bielory, L., Nguyen, V., Diaz, R. & Jeng, H.A.** 2012. Anthropogenic climate change and allergic diseases. Atmosphere 3: 200-212.
- Bortenschlager, S. & Bortenschlager, I. 2005. Altering airborne pollen concentrations due to the Global Warming. A comparative analysis of airborne pollen records from Innsbruck and Obergurgl (Austria) for the period 1980–2001. Grana 44: 172-180.
- **Branzi, G.P. & Zanotti, A.L.** 1992. Estimate and mapping of the activity of airborne pollen sources. Aerobiologia 8: 69-74.
- Bruffaerts, N., De Smedt, T., Delcloo, A., Simons, K., Hoebeke, L., Verstraeten, C., Van Nieuwenhuyse, A., Packeu, A. & Hendrickx, M. 2018. Comparative long-term trend analysis of daily weather conditions with daily pollen concentrations in Brussels, Belgium. International Journal of Biometeorology 62: 483-491.
- Burr, M.L. 1999. Grass pollen: trends and predictions. Clinical and experimental allergy: journal of the British Society for Allergy and Clinical Immunology 29: 735-738.
- Buters, J.T., Antunes, C., Galveias, A., Bergmann,
 K.C., Thibaudon, M., Galán, C., Schmidt-Weber,
 C. & Oteros, J. 2018. Pollen and spore monitoring in
 the world. Clinical and Translational Allergy 8: 9.
- Caillaud, D., Thibaudon, M., Martin, S., Ségala, C., Besancenot, J., Clot, B. & François, H. 2014. Shortterm effects of airborne ragweed pollen on clinical symptoms of hay fever in a panel of 30 patients. Journal of Investigational Allergology and Clinical Immunology 24: 249-256.
- Cariñanos González, P. & Casares Porcel, M. 2019. Estimation of the Allergenic Potential of Urban Trees and Urban Parks: Towards the Healthy Design of Urban Green Spaces of the Future.
- Cariñanos, P., Alcázar, P., Galán, C. & Domínguez, E. 2014. Environmental behaviour of airborne Amaranthaceae pollen in the southern part of the Iberian Peninsula, and its role in future climate scenarios. Science of the Total Environment 470: 480-487.
- Cariñanos, P. & Casares-Porcel, M. 2011. Urban green zones and related pollen allergy: A review. Some guidelines for designing spaces with low allergy impact. Landscape and Urban Planning 101: 205-214.
- Cecchi, L., D'amato, G., Ayres, J., Galan, C., Forastiere, F., Forsberg, B., Gerritsen, J., Nunes, C., Behrendt, H. & Akdis, C. 2010. Projections of the effects of climate change on allergic asthma: the contribution of aerobiology. Allergy 65: 1073-1081.
- Charpin, D., Pichot, C., Belmonte, J., Sutra, J.P., Zidkova, J., Chanez, P., Shahali, Y., Sénéchal, H. & Poncet, P. 2019. Cypress pollinosis: from tree to clinic. Clinical Reviews in Allergy & Immunology 56: 174-195.
- Chassard, G., Choel, M., Gosselin, S., Vorng, H., Petitprez, D., Shahali, Y., Tsicopoulos, A. & Visez, N. 2015. Kinetic of NO2 uptake by Phleum pratense

- pollen: chemical and allergenic implications. Environmental Pollution 196: 107-113.
- Chuine, I., Cour, P. & Rousseau, D. 1998. Fitting models predicting dates of flowering of temperate zone trees using simulated annealing. Plant, Cell & Environment 21: 455-466.
- **Clot, B.** 2003. Trends in airborne pollen: an overview of 21 years of data in Neuchâtel (Switzerland). Aerobiologia 19: 227-234.
- **Corden, J.M. & Millington, W.M.** 2001. The long-term trends and seasonal variation of the aeroallergen Alternaria in Derby, UK. Aerobiologia 17: 127-136.
- Cristofori, A., Cristofolini, F. & Gottardini, E. 2010. Twenty years of aerobiological monitoring in Trentino (Italy): assessment and evaluation of airborne pollen variability. Aerobiologia 26: 253-261.
- Curtis, P.S. & Wang, X. 1998. A meta-analysis of elevated CO 2 effects on woody plant mass, form, and physiology. Oecologia 113: 299-313.
- **D'amato, G.** 2011. Effects of climatic changes and urban air pollution on the rising trends of respiratory allergy and asthma. Multidisciplinary respiratory medicine 6: 28.
- D browska-Zapart, K., Chłopek, K. & Nied wied, T. 2018. The impact of meteorological conditions on the concentration of alder pollen in Sosnowiec (Poland) in the years 1997–2017. Aerobiologia 34: 469-485.
- Damialis, A., Gioulekas, D., Lazopoulou, C., Balafoutis, C. & Vokou, D. 2005. Transport of airborne pollen into the city of Thessaloniki: the effects of wind direction, speed and persistence. International Journal of Biometeorology 49: 139-145.
- Damialis, A., Halley, J.M., Gioulekas, D. & Vokou, D. 2007. Long-term trends in atmospheric pollen levels in the city of Thessaloniki, Greece. Atmospheric Environment 41: 7011-7021.
- De León, D.G., García-Mozo, H., Galán, C., Alcázar, P., Lima, M. & González-Andújar, J.L. 2015. Disentangling the effects of feedback structure and climate on Poaceae annual airborne pollen fluctuations and the possible consequences of climate change. Science of the Total Environment 530: 103-109.
- **Defila, C. & Clot, B.** 2001. Phytophenological trends in Switzerland. International Journal of Biometeorology 45: 203-207.
- Donders, T.H., Hagemans, K., Dekker, S.C., De Weger, L.A., De Klerk, P. & Wagner-Cremer, F. 2014. Region-specific sensitivity of anemophilous pollen deposition to temperature and precipitation. PloS One 9.
- **Doran, P.T. & Zimmerman, M.K.** 2009. Examining the scientific consensus on climate change. Eos, Transactions American Geophysical Union 90: 22-23.
- Duque, L., Guimarães, F., Ribeiro, H., Sousa, R. & Abreu, I. 2013. Elemental characterization of the airborne pollen surface using Electron Probe Microanalysis (EPMA). Atmospheric Environment 75: 296-302.
- Emberlin, J., Detandt, M., Gehrig, R., Jaeger, S., Nolard, N. & Rantio-Lehtimäki, A. 2002. Responses in the start of *Betula* (birch) pollen seasons

- to recent changes in spring temperatures across Europe. International Journal of Biometeorology 46: 159-170.
- Emberlin, J., Mullins, J., Corden, J., Jones, S., Millington, W., Brooke, M. & Savage, M. 1999. Regional variations in grass pollen seasons in the UK, long-term trends and forecast models. Clinical and experimental allergy: journal of the British Society for Allergy and Clinical Immunology 29: 347-356.
- Emberlin, J., Mullins, J., Corden, J., Millington, W., Brooke, M., Savage, M. & Jones, S. 1997. The trend to earlier birch pollen seasons in the UK: a biotic response to changes in weather conditions? Grana 36: 29-33.
- Emberlin, J. & Norris-Hill, J. 1991. Spatial variation of pollen deposition in North London. Grana 30: 190-195.
- **Fitter, A. & Fitter, R.** 2002. Rapid changes in flowering time in British plants. Science 296: 1689-1691.
- Frank, U. & Ernst, D. 2016. Effects of NO2 and ozone on pollen allergenicity. Frontiers in Plant Science 7: 91.
- **Frei, T. & Gassner, E.** 2008. Climate change and its impact on birch pollen quantities and the start of the pollen season an example from Switzerland for the period 1969–2006. International Journal of Biometeorology 52: 667.
- **Frenguelli, G.** 2002. Interactions between climatic changes and allergenic plants. Monaldi Archives for Chest Disease 57: 141-143.
- Galán, C., Alcázar, P., Oteros, J., García-Mozo, H., Aira, M., Belmonte, J., De La Guardia, C.D., Fernández-González, D., Gutierrez-Bustillo, M. & Moreno-Grau, S. 2016. Airborne pollen trends in the Iberian Peninsula. Science of the Total Environment 550: 53-59.
- Galán, C., Ariatti, A., Bonini, M., Clot, B., Crouzy, B., Dahl, A., Fernandez-González, D., Frenguelli, G., Gehrig, R. & Isard, S. 2017. Recommended terminology for aerobiological studies. Aerobiologia 33: 293-295.
- Galán, C., García-Mozo, H., Vázquez, L., Ruiz, L., De La Guardia, C.D. & Trigo, M. 2005. Heat requirement for the onset of the *Olea europaea* L. pollen season in several sites in Andalusia and the effect of the expected future climate change. International Journal of Biometeorology 49: 184-188.
- **Gallego, M., García, J. & Vaquero, J.** 2005. The NAO signal in daily rainfall series over the Iberian Peninsula. Climate Research 29: 103-109.
- García-Mozo, H., Galán, C., Belmonte, J., Bermejo, D., Candau, P., De La Guardia, C.D., Elvira, B., Gutiérrez, M., Jato, V. & Silva, I. 2009. Predicting the start and peak dates of the Poaceae pollen season in Spain using process-based models. Agricultural and Forest Meteorology 149: 256-262.
- García-Mozo, H., Galán, C., Jato, V., Belmonte, J., De La Guardia, C.D., Fernández, D., Gutiérrez, M., Aira, M.J., Roure, J.M. & Ruiz, L. 2006. *Quercus* pollen season dynamics in the Iberian Peninsula: response to meteorological parameters and possible consequences of climate change. Annals of

- Agricultural and Environmental Medicine 13: 209.
- Garcia-Mozo, H., Orlandi, F., Galan, C., Fornaciari, M., Romano, B., Ruiz, L., De La Guardia, C.D., Trigo, M. & Chuine, I. 2009. Olive flowering phenology variation between different cultivars in Spain and Italy: modeling analysis. Theoretical and Applied Climatology 95: 385.
- García-Mozo, H., Oteros, J.A. & Galán, C. 2016. Impact of land cover changes and climate on the main airborne pollen types in Southern Spain. Science of the Total Environment 548: 221-228.
- García-Mozo, H., Yaezel, L., Oteros, J. & Galán, C. 2014. Statistical approach to the analysis of olive long-term pollen season trends in southern Spain. Science of the Total Environment 473: 103-109.
- Georgakopoulos, D., Després, V., Fröhlich-Nowoisky, J., Psenner, R., Ariya, P., Pósfai, M., Ahern, H., Moffett, B. & Hill, T. 2008. Microbiology and atmospheric processes: biological, physical and chemical characterization of aerosol particles.
- Ghahremaninejad, F., Hoseini, E. & Fereidounfar, S. 2021. Cities in drylands as artificial protected areas for plants. Biodiversity and Conservation 30: 243-248
- Girgis, S., Marks, G., Downs, S., Kolbe, A., Car, G. & Paton, R. 2000. Thunderstorm-associated asthma in an inland town in south-eastern Australia. Who is at risk? European Respiratory Journal 16: 3-8.
- Green, B. J., Dettmann, M., Yli-Panula, E., Rutherford, S. & Simpson, R. 2004. Atmospheric Poaceae pollen frequencies and associations with meteorological parameters in Brisbane, Australia: a 5-year record, 1994–1999. International Journal of Biometeorology 48: 172-178.
- Häfner, D., Reich, K., Matricardi, P., Meyer, H., Kettner, J. & Narkus, A. 2011. Prospective validation of 'Allergy Control SCORETM': a novel symptom-medication score for clinical trials. Allergy 66: 629-636.
- Hamaoui-Laguel, L., Vautard, R., Liu, L., Solmon, F., Viovy, N., Khvorostyanov, D., Essl, F., Chuine, I., Colette, A. & Semenov, M.A. 2015. Effects of climate change and seed dispersal on airborne ragweed pollen loads in Europe. Nature Climate Change 5: 766-771.
- Hernández-Ceballos, M., García-Mozo, H., Adame, J., Domínguez-Vilches, E., Bolívar, J., De La Morena, B., Pérez-Badía, R. & Galán, C. 2011. Determination of potential sources of *Quercus* airborne pollen in Córdoba city (southern Spain) using back-trajectory analysis. Aerobiologia 27: 261-276.
- **Hirst, J.M.** 1952. An automatic volumetric spore trap. Annals of applied Biology 39: 257-265.
- Huffman, J.A., Prenni, A., Demott, P., Pöhlker, C., Mason, R., Robinson, N., Fröhlich-Nowoisky, J., Tobo, Y., Després, V. & Garcia, E. 2013. High concentrations of biological aerosol particles and ice nuclei during and after rain. Atmospheric Chemistry and Physics 13: 6151.
- **Huynen, M.** 2003. Phenology and Human Health: Allergic Disorders. Report of a WHO meeting Rome,

- Italy. 16–17 January 2003. World Health Organization, Regional Office for Europe.
- Jablonski, L.M., Wang, X. & Curtis, P.S. 2002. Plant reproduction under elevated CO2 conditions: a meta analysis of reports on 79 crop and wild species. New Phytologist 156: 9-26.
- Jäger, S., Spieksma, E.T.M. & Nolard, N. 1991. Fluctuations and trends in airborne concentrations of some abundant pollen types, monitored at Vienna, Leiden, and Brussels. Grana 30: 309-312.
- Jato, V., Rodríguez-Rajo, F., Seijo, M. & Aira, M. 2009. Poaceae pollen in Galicia (NW Spain): characterisation and recent trends in atmospheric pollen season. International Journal of Biometeorology 53: 333.
- Jochner, S., Lüpke, M., Laube, J., Weichenmeier, I., Pusch, G., Traidl-Hoffmann, C., Schmidt-Weber, C., Buters, J.T. & Menzel, A. 2015. Seasonal variation of birch and grass pollen loads and allergen release at two sites in the German Alps. Atmospheric Environment 122: 83-93.
- José González Minero, F. & Candau, P. 1997. Study on Platanus hispanica Miller pollen content in the air of Seville, southern Spain. Aerobiologia 13: 109-115.
- Katotomichelakis, M., Nikolaidis, C., Makris, M., Zhang, N., Aggelides, X., Constantinidis, T.C., Bachert, C. & Danielides, V. (2015). The clinical significance of the pollen calendar of the Western Thrace/northeast Greece region in allergic rhinitis. Paper presented at the International forum of allergy & rhinology.
- Kazinczi, G., Novák, R., Pathy, Z. & Béres, I. 2008. Common ragweed (*Ambrosia artemisiifolia* L.): a review with special regards to the results in Hungary. III. Resistant biotypes, control methods and authority arrangements. Herbologia 9: 119-144.
- Kitinoja, M.A., Hugg, T.T., Siddika, N., Yanez, D.R., Jaakkola, M.S. & Jaakkola, J.J. 2020. Short-term exposure to pollen and the risk of allergic and asthmatic manifestations: a systematic review and meta-analysis. BMJ Open 10.
- Knapp, P.A. & Soulé, P.T. 1998. Recent Juniperus occidentalis (western juniper) expansion on a protected site in central Oregon. Global Change Biology 4: 347-357.
- **Ladeau, S.L. & Clark, J.** 2006. Pollen production by *Pinus taeda* growing in elevated atmospheric CO2. Functional Ecology 541-547.
- **Lavee, S. & Datt, A.** 1978. The necessity of cross-pollination for fruit set of Manzanillo olives. Journal of Horticultural Science 53: 261-266.
- Linderholm, H.W. 2006. Growing season changes in the last century. Agricultural and Forest Meteorology 137: 1-14.
- Lo, F., Bitz, C.M. Battisti, D.S., & Hess, J.J. 2019. Pollen calendars and maps of allergenic pollen in North America. Aerobiologia 35: 613-633.
- Mansouritorghabeh, H., Jabbari-Azad, F., Sankian, M., Varasteh, A. & Farid-Hosseini, R. 2019. The most common allergenic tree pollen grains in the Middle East: a narrative review. Iranian Journal of Medical Sciences 44: 87.

- **Menzel, A.** 2000. Trends in phenological phases in Europe between 1951 and 1996. International Journal of Biometeorology 44: 76-81.
- **Menzel, A. & Fabian, P.** 1999. Growing season extended in Europe. Nature 397: 659-659.
- Mimet, A., Pellissier, V., Quénol, H., Aguejdad, R., Dubreuil, V. & Roze, F. 2009. Urbanisation induces early flowering: evidence from *Platanus acerifolia* and *Prunus cerasus*. International Journal of Biometeorology 53: 287-298.
- Minero, F.G., Candau, P., Tomas, C. & Morales, J. 1998. Airborne grass (Poaceae) pollen in southern Spain. Results of a 10 year study (1987–96). Allergy 53: 266-274.
- **Moore, L.M. & Lauenroth, W.K.** 2017. Differential effects of temperature and precipitation on early vs. late flowering species. Ecosphere 8: 01819.
- Mousavi, F., Majd, A., Shahali, Y., Ghahremaninejad, F., Kardar, G. & Pourpak, Z. 2016. Pollinosis to tree of heaven (*Ailanthus altissima*) and detection of allergenic proteins: a case report. Annals of Allergy, Asthma & Immunology 116: 374-375.
- Mousavi, F., Shahali, Y., Pourpak, Z., Majd, A. & Ghahremaninejad, F. 2019. Year-to-year variation of the elemental and allergenic contents of *Ailanthus altissima* pollen grains: an allergomic study. Environmental Monitoring and Assessment 191: 362.
- Müller-Germann, I., Vogel, B., Vogel, H., Pauling, A., Fröhlich-Nowoisky, J., Pöschl, U. & Després, V.R. 2015. Quantitative DNA analyses for airborne birch pollen. PloS One 10.
- Norby, R.J., Wullschleger, S.D., Gunderson, C.A., Johnson, D.W. & Ceulemans, R. 1999. Tree responses to rising CO2 in field experiments: implications for the future forest. Plant, Cell & Environment 22: 683-714.
- Oduber, F., Calvo, A., Blanco-Alegre, C., Castro, A., Vega-Maray, A., Valencia-Barrera, R., Fernández-González, D. & Fraile, R. 2019. Links between recent trends in airborne pollen concentration, meteorological parameters and air pollutants. Agricultural and Forest Meteorology 264: 16-26.
- Okuyama, Y., Matsumoto, K., Okochi, H. & Igawa, M. 2007. Adsorption of air pollutants on the grain surface of Japanese cedar pollen. Atmospheric Environment 41: 253-260.
- Ørby, P.V., Peel, R.G., Skjøth, C., Schlünssen, V., Bønløkke, J., Ellermann, T., Brændholt, A., Sigsgaard, T. & Hertel, O. 2015. An assessment of the potential for co-exposure to allergenic pollen and air pollution in Copenhagen, Denmark. Urban Climate 14: 457-474.
- Orlandi, F., Garcia-Mozo, H., Galán, C., Romano, B., De La Guardia, C.D., Ruiz, L., Del Mar Trigo, M., Dominguez-Vilches, E. & Fornaciari, M. 2010. Olive flowering trends in a large Mediterranean area (Italy and Spain). International Journal of Biometeorology 54: 151-163.
- Oteros, J., García-Mozo, H., Hervás-Martínez, C. & Galán, C. 2013a. Year clustering analysis for modelling olive flowering phenology. International Journal of Biometeorology 57: 545-555.

- Oteros, J., García-Mozo, H., Hervás, C. & Galán, C. 2013b. Biometeorological and autoregressive indices for predicting olive pollen intensity. International Journal of Biometeorology 57: 307-316.
- Oteros, J., García-Mozo, H., Vázquez, L., Mestre, A., Domínguez-Vilches, E. & Galán, C. 2013c. Modelling olive phenological response to weather and topography. Agriculture, Ecosystems & Environment 179: 62-68.
- Palacios, I. S., Molina, R.T. & Rodríguez, A.M. 2000. Influence of wind direction on pollen concentration in the atmosphere. International Journal of Biometeorology 44: 128-133.
- Pawankar, R., Canonica, G., Holgate, S., Lockey, R. & Blaiss, M. 2011. World Allergy Organization (WAO) white book on allergy. Wisconsin: World Allergy Organisation.
- Pfaar, O., Bastl, K., Berger, U., Buters, J., Calderon, M., Clot, B., Darsow, U., Demoly, P., Durham, S. & Galán, C. 2017. Defining pollen exposure times for clinical trials of allergen immunotherapy for pollen induced rhinoconjunctivitis—an EAACI position paper. Allergy 72: 713-722.
- Puc, M. 2012. Artificial neural network model of the relationship between *Betula* pollen and meteorological factors in Szczecin (Poland). International Journal of Biometeorology 56: 395-401.
- **Rasmussen, A.** 2002. The effects of climate change on the birch pollen season in Denmark. Aerobiologia 18: 253-265.
- Ravindra, K., Goyal, A., Kumar, S., Aggarwal, A. & Mor, S. 2021. Pollen Calendar to depict seasonal periodicities of airborne pollen species in a city situated in Indo-Gangetic plain, India. Atmospheric Environment 262: 118649.
- Recio, M., Picornell, A., Trigo, M., Gharbi, D., García-Sánchez, J. & Cabezudo, B. 2018. Intensity and temporality of airborne *Quercus* pollen in the southwest Mediterranean area: Correlation with meteorological and phenoclimatic variables, trends and possible adaptation to climate change. Agricultural and Forest Meteorology 250: 308-318.
- Recio, M., Rodríguez-Rajo, F.J., Jato, M.V., Trigo, M.M. & Cabezudo, B. 2009. The effect of recent climatic trends on Urticaceae pollination in two bioclimatically different areas in the Iberian Peninsula: Malaga and Vigo. Climatic Change 97: 215-228.
- Ridolo, E., Albertini, R., Giordano, D., Soliani, L., Usberti, I. & Dall'aglio, P. 2007. Airborne pollen concentrations and the incidence of allergic asthma and rhinoconjunctivitis in northern Italy from 1992 to 2003. International Archives of Allergy and Immunology 142: 151-157.
- Roberts, G., Mylonopoulou, M., Hurley, C. & Lack, G. 2005. Impairment in quality of life is directly related to the level of allergen exposure and allergic airway inflammation. Clinical & Experimental Allergy 35: 1295-1300.
- Rodriguez-Rajo, F.J., Dopazo, A. & Jato, V. 2004. Environmental factors affecting the start of pollen season and concentrations of airborne Alnus pollen in

- two localities of Galicia (NW Spain). Annals of Agricultural and Environmental Medicine 11: 35-44.
- Rogers, C.A., Wayne, P.M., Macklin, E.A., Muilenberg, M.L., Wagner, C.J., Epstein, P.R. & Bazzaz, F.A. 2006. Interaction of the onset of spring and elevated atmospheric CO2 on ragweed (*Ambrosia artemisiifolia* L.) pollen production. Environmental Health Perspectives 114: 865-869.
- **Rogoff, M.J.** 2013. Solid waste recycling and processing: planning of solid waste recycling facilities and programs: Elsevier.
- Rojo, J., Orlandi, F., Ben Dhiab, A., Lara, B., Picornell, A., Oteros, J., Msallem, M., Fornaciari, M. & Pérez-Badia, R. 2020. Estimation of Chilling and Heat Accumulation Periods Based on the Timing of Olive Pollination. Forests 11: 835.
- **Rojo, J. & Pérez-Badia, R.** 2015. Spatiotemporal analysis of olive flowering using geostatistical techniques. Science of the Total Environment 505: 860-869.
- Rojo, J., Rapp, A., Lara, B., Fernández-González, F. & Pérez-Badia, R. 2015. Effect of land uses and wind direction on the contribution of local sources to airborne pollen. Science of the Total environment 538: 672-682.
- Romero-Morte, J., Rojo, J. & Pérez-Badia, R. 2020. Meteorological factors driving airborne grass pollen concentration in central Iberian Peninsula. Aerobiologia 36: 527-540.
- Ruiz-Valenzuela, L. & Aguilera, F. 2018. Trends in airborne pollen and pollen-season-related features of anemophilous species in Jaen (south Spain): A 23year perspective. Atmospheric Environment 180: 234-243.
- Sauliene, I., Sukiene, L. & Kazlauskiene, V. 2019. The assessment of atmospheric conditions and constituents on allergenic pollen loads in Lithuania. Journal of Environmental Management 250: 109469.
- Š evková, J., Duši ka, J., Hrabovský, M. & Vašková, Z. 2021. Trends in pollen season characteristics of Alnus, Poaceae and *Artemisia* allergenic taxa in Bratislava, central Europe. Aerobiologia1-11.
- Schäppi, G.F., Suphioglu, C., Taylor, P.E. & Knox, R.B. 1997. Concentrations of the major birch tree allergen Bet v 1 in pollen and respirable fine particles in the atmosphere. Journal of Allergy and Clinical Immunology 100: 656-661.
- Scheifinger, H., Belmonte, J., Buters, J., Celenk, S., Damialis, A., Dechamp, C., García-Mozo, H., Gehrig, R., Grewling, L. & Halley, J.M. (2013). Monitoring, Modelling and Forecasting of the pollen season Allergenic pollen (pp. 71-126): Springer.
- **Schwartz, M.D.** 2003. Phenology: an integrative environmental science: Springer.
- Sénéchal, H., Visez, N., Charpin, D., Shahali, Y., Peltre, G., Biolley, J.P., Lhuissier, F., Couderc, R., Yamada, O. & Malrat-Domenge, A. 2015. A review of the effects of major atmospheric pollutants on pollen grains, pollen content, and allergenicity. The Scientific World Journal 2015.
- Shahali, Y., Pourpak, Z., Moin, M., Zare, A. & Majd, A. 2009. Impacts of air pollution exposure on the

- allergenic properties of Arizona cypress pollens. Paper presented at the Journal of Physics: Conference Series.
- Sherry, R.A., Zhou, X., Gu, S., Arnone, J.A., Schimel, D.S., Verburg, P.S., Wallace, L.L. & Luo, Y. 2007. Divergence of reproductive phenology under climate warming. Proceedings of the National Academy of Sciences 104: 198-202.
- Singer, B.D., Ziska, L.H., Frenz, D.A., Gebhard, D.E. & Straka, J.G. 2005. Increasing Amb a 1 content in common ragweed (*Ambrosia artemisiifolia*) pollen as a function of rising atmospheric CO2 concentration. Functional Plant Biology 32: 667-670.
- Smith, M., Skjøth, C.A., Myszkowska, D., Uruska, A., Puc, M., Stach, A., Balwierz, Z., Chlopek, K., Piotrowska, K. & Kasprzyk, I. 2008. Long-range transport of *Ambrosia* pollen to Poland. Agricultural and Forest Meteorology 148: 1402-1411.
- Sofiev, M., Belmonte, J., Gehrig, R., Izquierdo, R., Smith, M., Dahl, Å. & Siljamo, P. (2013). Airborne pollen transport Allergenic pollen (pp. 127-159): Springer.
- Solomon, S., Manning, M., Marquis, M. & Qin, D. 2007. Climate change 2007-the physical science basis: Working group I contribution to the fourth assessment report of the IPCC (Vol. 4): Cambridge university press.
- Spieksma, F.T.M., Corden, J., Detandt, M., Millington, W., Nikkels, H., Nolard, N., Schoenmakers, C., Wachter, R., De Weger, L. & Willems, R. 2003. Quantitative trends in annual totals of five common airborne pollen types (*Betula*, *Quercus*, Poaceae, *Urtica*, and *Artemisia*), at five pollen-monitoring stations in western Europe. Aerobiologia 19: 171-184.
- Spieksma, F.T.M., Emberlin, J., Hjelmroos, M., Jäger, S. & Leuschner, R. 1995. Atmospheric birch (*Betula*) pollen in Europe: trends and fluctuations in annual quantities and the starting dates of the seasons. Grana 34: 51-57.
- Srikanth, P., Sudharsanam, S. & Steinberg, R. 2008. Bio-aerosols in indoor environment: composition, health effects and analysis. Indian Journal of Medical Microbiology 26: 302.
- Stach, A., García-Mozo, H., Prieto-Baena, J., Czarnecka-Operacz, M., Jenerowicz, D., Silny, W. & Galán, C. 2007. Prevalence of Artemisia species pollinosis in western Poland: Impact of climate change on aerobiological trends. J Investig Allergol Clin Immunol 17: 39-47.
- Suárez-Varela, M.M., Alvarez, L. G.M., Kogan, M.D., González, A.L., Gimeno, A.M., Ontoso, I.A., Díaz, C.G., Pena, A.A., Aurrecoechea, B.D. & Monge, R.M.B. 2008. Climate and prevalence of atopic eczema in 6-to 7-year-old school children in Spain. ISAAC phase III. International Journal of Biometeorology 52: 833-840.
- Taylor, P.E., Jacobson, K.W., House, J.M. & Glovsky, M.M. 2007. Links between pollen, atopy and the asthma epidemic. International Archives of Allergy and Immunology 144: 162-170.
- Tedeschini, E., Javier Rodríguez Rajo, F.,

- Caramiello, R., Jato, V. & Frenguelli, G. 2006. The influence of climate changes in *Platanus* spp. pollination in Spain and Italy. Grana 45: 222-229.
- Thien, F., Beggs, P.J., Csutoros, D., Darvall, J., Hew, M., Davies, J.M., Bardin, P.G., Bannister, T., Barnes, S. & Bellomo, R. 2018. The Melbourne epidemic thunderstorm asthma event 2016: an investigation of environmental triggers, effect on health services, and patient risk factors. The Lancet Planetary Health 2: 255-263.
- Tosunoglu, A., Altunoglu, M.K., Bicakci, A., Kilic, O., Gonca, T., Yilmazer, I., Saatcioglu, G., Akkaya, A., Celenk, S. & Canitez, Y. 2015. Atmospheric pollen concentrations in Antalya, South Turkey. Aerobiologia 31: 99-109.
- Uguz, U., Guvensen, A. & Tort, N.S. 2017. Annual and intradiurnal variation of dominant airborne pollen and the effects of meteorological factors in Çe me (Izmir, Turkey). Environmental Monitoring and Assessment 189: 530.
- **Unger, J.** 1999. Comparisons of urban and rural bioclimatological conditions in the case of a Central-European city. International Journal of Biometeorology 43: 139-144.
- Ursu C, I.C. Veres L. 2012. Management of respirato¬ries allergies in the context of sustainable development pro¬grams., from Alergologia http://www.sraic.eu/inpage/alergolo¬gia-1-2012/ (abstract)
- Van Vliet, A.J., Overeem, A., De Groot, R.S., Jacobs, A.F. & Spieksma, F.T. 2002. The influence of temperature and climate change on the timing of pollen release in the Netherlands. International Journal of Climatology: A Journal of the Royal Meteorological Society 22: 1757-1767.
- Voltolini, S., Minale, P., Troise, C., Bignardi, D., Modena, P., Arobba, D. & Negrini, A.C. 2000. Trend of herbaceous pollen diffusion and allergic sensitisation in Genoa, Italy. Aerobiologia 16: 245-249.
- Walther, G.R., Post, E., Convey, P., Menzel, A., Parmesan, C., Beebee, T.J., Fromentin, J.M., Hoegh-Guldberg, O. & Bairlein, F. 2002. Ecological responses to recent climate change. Nature 416: 389.
- Wan, S., Yuan, T., Bowdish, S., Wallace, L., Russell, S.D. & Luo, Y. 2002. Response of an allergenic species, *Ambrosia psilostachya* (Asteraceae), to experimental warming and clipping: implications for public health. American Journal of Botany 89: 1843-1846.
- Wayne, P., Foster, S., Connolly, J., Bazzaz, F. & Epstein, P. 2002. Production of allergenic pollen by ragweed (*Ambrosia artemisiifolia* L.) is increased in CO2-enriched atmospheres. Annals of Allergy, Asthma & Immunology 88: 279-282.
- Yli-Panula, E., Fekedulegn, D.B., Green, B.J. & Ranta, H. 2009. Analysis of airborne *Betula* pollen in Finland; a 31-year perspective. International journal of environmental research and public health 6: 1706-1723
- Zhang, Y., Bielory, L., Mi, Z., Cai, T., Robock, A. &

[DOI: 10.52547/nbr.9.2.95]

- **Georgopoulos, P.** 2015. Allergenic pollen season variations in the past two decades under changing climate in the United States. Global Change Biology 21: 1581-1589.
- **Ziska, L., George, K. & Frenz, D.** 2007. Establishment and persistence of common ragweed (*Ambrosia artemisiifolia* L.) in disturbed soil as a function of an urban–rural macro environment. Global Change Biology 13: 266-274.
- Ziska, L.H. & Beggs, P.J. 2012. Anthropogenic climate change and allergen exposure: the role of plant biology. Journal of Allergy and Clinical Immunology 129: 27-32.
- **Ziska, L.H. & Caulfield, F.A.** 2000. Rising CO2 and pollen production of common ragweed (*Ambrosia*

- artemisiifolia L.), a known allergy-inducing species: implications for public health. Functional Plant Biology 27: 893-898.
- Ziska, L.H. Gebhard, D.E., Frenz, D.A., Faulkner, S., Singer, B.D., & Straka, J.G. 2003. Cities as harbingers of climate change: common ragweed, urbanization, and public health. Journal of Allergy and Clinical Immunology 111: 290-295.
- Ziska, L.H. Makra, L., Harry, S.K., Bruffaerts, N., Hendrickx, M., Coates, F., Saarto, A., Thibaudon, M., Oliver, G., & Damialis, A. 2019. Temperature-related changes in airborne allergenic pollen abundance and seasonality across the northern hemisphere: a retrospective data analysis. The Lancet Planetary Health 3: 124-131.

How to cite this article

Mousavi, F., Shahali, Y., Oteros, J. & Bergmann, K-Ch. 2022. The impacts of climate change and meteorological factors on pollen season indicators of allergenic plant taxa. Nova Biologica Reperta 9: 95-114.